

arXiv:1510.06738v1:

Cold-gas outflows in typical low-redshift galaxies are driven by star formation, not AGN

Marc Sarzi^{1*}, Sugata Kaviraj¹, Borislav Nedelchev¹, Joshua Tiffany², Stanislav S. Shabala³, Adam T. Deller⁴ and Enno Middelberg⁵

¹*Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield AL10 9AB, United Kingdom*

²*Astronomy Unit, School of Physics and Astronomy, Queen Mary University of London, London E1 4NS, UK*

³*School of Mathematics and Physics, University of Tasmania, Private Bag 37, Hobart, TAS 7001, Australia*

⁴*The Netherlands Institute for Radio Astronomy (ASTRON), Dwingeloo, The Netherlands*

⁵*Astronomisches Institut der Ruhr-Universität Bochum, Universitätsstrae 150, D-44801 Bochum, Germany*



- mJIVE-20 (VLBI: $>1.2\text{mJy}$, seeing $0.01''$) + value-added SDSS catalogue OSSY (Oh et al. 2011)
 - 456 targets, 10^{11}Msun , 58% - ETG, 39% - LTG, 3% - mergers
 - 23% (103 galaxies) have VLBI detections, 90% of which occur in ETGs
- Rupke, Veilleux, & Sanders (2005) approach for NaD



